

Graphical Abstract

To create your abstract, type over the instructions in the template box below.
Fonts or abstract dimensions should not be changed or altered.

Preparation of Siloxacyclopentene Containing 1,3-Dienes and Their Diels-Alder Reactions

Leave this area blank for abstract info.



R = Me (88%, *endo*)
= ^tPr (86%, -1:1, *exo to endo*)

Ramakrishna R. Pidaparathi and Mark E. Welker*

Department of Chemistry, Wake Forest University, P.O Box 7486, Winston-Salem, NC 27109, USA

A number of enynyloxy dimethyl and diisopropyl silanes have been prepared and converted into siloxacyclopentene containing 1,3 dienes via intramolecular hydrosilylation of the alkyne functional group. Diels-Alder reactions of these dienes are reported.



Pergamon

TETRAHEDRON
LETTERS

Preparation of Siloxacyclopentene Containing 1,3-Dienes and Their Diels-Alder Reactions

Ramakrishna R. Pidaparathi and Mark E. Welker*

Department of Chemistry, Wake Forest University, P.O. Box 7486, Winston-Salem, NC 27109, USA.


Abstract— A number of enynyloxy dimethyl and diisopropyl silanes have been prepared and converted into siloxacyclopentene containing 1,3-dienes via intramolecular hydrosilylation of the alkyne functional group. Diels-Alder reactions of these dienes are reported. © 2011 Elsevier Science. All rights reserved

We have embarked on a synthetic program aimed at the synthesis of stable 2-boron and 2-silicon substituted 1,3-dienes and investigation of their reaction chemistry.^{1,2} Earlier this year we reported the simple preparation of 2-triethoxysilyl-1,3-butadiene and its conversion into 2-silatrane and 2-biscatecholsiloxy-substituted 1,3 dienes. Those new siloxy substituted dienes proved to be remarkably reactive in Diels-Alder cycloaddition reactions and the Diels-Alder cycloadducts could be used in Hiyama cross coupling reactions.² Reports of the preparation and use of 2-trialkoxysilyl-1,3-dienes are extremely rare. We find a report of 2-trimethoxysilyl- and 2-triethoxysilyl-1,3-butadiene in 1984³ and then a report of the polymerization of these materials in 1989.⁴

In 1988, Tamao, Ito and co-workers reported the intramolecular hydrosilylation of the alkyne moiety of homopropargyl alcohols to make siloxacyclopentanes containing exocyclic alkenes.⁵ Other routes to siloxacycle containing species which have been used in cross coupling

chemistry containing a silicon substituent at the 1 position of a 1,3 diene moiety.⁹ This diene reacted with N-phenylmaleimide in a Diels-Alder reaction. Also in 2004 Lee and co-workers reported the synthesis of a number of siloxacycles that are part of a 1,3-diene unit via a condensation/metathesis strategy using alkenyl alcohols and alkynyl silanes.¹⁰ No Diels-Alder or cross coupling reactions of these substrates were reported. Earlier this year Halvorsen and Roush reported using this procedure to make siloxacyclopentenes containing pendant dienophiles and demonstrated that they could be used in intramolecular Diels-Alder reactions which were followed by protodesilylation.¹¹ We had simultaneously been using a combination of the Tamao-Ito⁵ and Lee protocols¹² to make siloxacyclopentenes which could participate in intermolecular Diels-Alder reactions. Here, we report our preliminary results in this area.

Scheme 1. Synthesis of Enynylsilanes (1-3).



chemistry

have also been reported previously.⁶⁻⁸ In 2004, Clark and Woerpel reported the reaction of a "tBu₂Si" source with a protected enynol to produce a siloxacyclopentene

enynyl silanes (Scheme 1) in excellent yields from pentenyne and tried to convert them into siloxacyclopentene containing 1,3 dienes (**4**) using Lee's


We first prepared several

* Corresponding author. Tel.: +1-336-758-5758; fax: +1-336-758-4321; e-mail: welker@wfu.edu

alkynylation-hydrosilylation sequence.¹² Unfortunately, we recovered enynyl silanes (**1-3**) from these reactions. We then switched to the preparation of siloxy substituted enynes (**7a, 7c-e**) (prepared from methylbutenyne (**5a**) and cyclohexenylethyne (**5b**) via condensation with carbonyl compounds followed by Denmark's protocol¹³ and

converted them into siloxacyclopentenes (**8a, 8e, 10a, 10e**) using Lee's KO^tBu catalyzed trans-hydrosilylation protocol¹² (Scheme 2). Diisopropylsilyloxy substituted enynes (**7a, 7c-e**) could be isolated and characterized prior to conversion to siloxacyclopentenes (**8a, 8e**).


Scheme 2. Synthesis of Diisopropylsilyloxy Substituted Enynes (**7a-e**) and Siloxacyclopentene Containing 1,3-Dienes (**8a, 8e**).



Conditions: a) i. THF, -78 °C; ii. ^tBuLi; iii. -78 °C → 0 °C, 1 h; b) i. 0 °C → -78 °C; ii. aldehyde/ketone; iii. -78 °C → r.t., overnight; c) i. hexanes, 0 °C; ii. DMAP, NEt₃; iii. ⁱPr₂Si(H)Cl in hex, 10 min; iv. 0 °C r.t., overnight; d) i. THF; ii. waterbath; iii. KO^tBu in THF; iv. 1 h.

Dimethylsilyloxy substituted enynes (**9a, 9e**) were generated in-situ by using the Tamo-Ito protocol⁵ and then converted into siloxacyclopentenes (**10a, 10e**) which were isolated


and characterized (Scheme 3). All dienes are stable for weeks at 25 °C in the presence of air and can be purified by chromatography on silica gel.

Scheme 3. Synthesis (Tandem) of Dimethyl Siloxacyclopentene Containing 1,3-Dienes (**10a**, **10e**).


Conditions: a) i. waterbath; ii. $(\text{Me}_2\text{SiH})_2\text{NH}$ (neat); iii. r.t., overnight; b) i. THF; ii. waterbath; iii. KO^tBu in THF; iv. r.t., 1 h.

With these dienes in hand we have performed several Diels-Alder reactions. Methyl substituted dienes **8a** and **10a** both react completely with N-phenylmaleimide after 24-36 h at 90 °C (Scheme 4). The dimethylsiloxy substituted diene (**10a**) provides *endo* adduct (**12**) whereas the more bulky diisopropyl siloxy diene **8a** provides *exo* and *endo* cycloadducts (**11a** and **11b** respectively) in almost equal proportions. This outcome can be rationalized by unfavorable steric interactions between the N-phenyl and the isopropyl groups in the electronically favored *endo* transition state. Cyclohexenyl substituted dienes **8e** and **10e** proved much less reactive in thermal Diels-Alder

reactions and only start to show traces of Diels-Alder cycloadducts after 40-50 h of heating at 90 °C. Halvorsen and Roush found that siloxacyclopentene constrained nonatrienes participated in thermal Diels-Alder reactions with little stereoselectivity whereas these substrates participated in Lewis acid catalyzed intramolecular Diels-Alder reactions through *endo* transition states to produce perhydroindene cycloadducts with high stereoselectivity.¹¹ They found that siloxacyclopentene constrained decatrienes participated in thermal or Lewis acid catalyzed cycloadditions through *endo* transition states to produce octahydronaphthalenes with high stereoselectivity.

Scheme 4. Intermolecular Diels-Alder Reactions of Diene - **8a** and **10a** with N-Phenylmaleimide.

The stereochemistry of the Diels-Alder cycloadducts **11** and **12** were assigned using a combination of COSY, HMQC, and HMBC to make all the ¹H assignments

Scheme 5. Cycloadduct Stereochemistry.

Endo adduct (**11b**) in C6D6 (other protons were not shown for clarity)

followed by NOESY to assign stereochemistry. In the *cis* isomers that arise from the *endo* transition states (**11b**, **12**),

one of the diastereotopic H5 protons exhibits NOE to both H8a and H8b whereas the other H5 does not exhibit NOE to either of those protons (Scheme 5).

In summary, we have prepared new diisopropylsiloxacyclopentene containing 1,3 dienes via a simple three step procedure from enynes. We have prepared new dimethylsiloxy cyclopentene containing 1,3 dienes via a one pot, two step sequence from enynols. These dienes participate in Diels-Alder reactions with a higher preference for the production of *trans* cycloadducts from the sterically more bulky diisopropylsiloxy substituted dienes.

Acknowledgement

We thank the National Science Foundation for their support of this work (CHE-0450722) and the NMR instrumentation used to characterize the compounds reported here. The Duke University Center for Mass Spectrometry and Mass used to characterize the compounds reported here. The Duke University Center for Mass Spectrometry and Mass Spectrometry Facility at UNC-CH performed high-resolution mass spectral analysis. We thank Dr. Marcus W. Wright for NMR assistance.

References and notes

- De, S.; Welker, M. E. *Org. Lett.* **2005**, *7*, 2481.
- Pidaparthy, R. R.; Welker, M. E.; Day, C. S.; Wright, M. W. *Org. Lett.* **2007**, *9*, 1623.
- Takenaka, K.; Hattori, T.; Hirao, A.; Nakahama, S. *Macromolecules*, **1989**, *22*, 1563.
- Sato, F.; Uchiyama, H.; Samaddar, A. K. *Chemistry & Industry*, **1984**, *20*, 743.
- Tamao, K.; Maeda, K.; Tanaka, T.; Ito, Y. *Tetrahedron Lett.* **1988**, *29*, 6955.
- Zacuto, M. J.; O'Malley, S. J.; Leighton, J. L. *J. Am. Chem. Soc.* **2002**, *124*, 7890.
- Trost, B. M.; Ball, Z. T.; Laemmerhold, K. M. *J. Am. Chem. Soc.* **2005**, *127*, 10028.
- Kim, Y. J.; Lee, D. *Org. Lett.* **2006**, *8*, 5219.
- Clark, T. B.; Woerpel, K. A. *J. Am. Chem. Soc.* **2004**, *126*, 9522.
- Miller, R. L.; Maifeld, S. V.; Lee, D. *Org. Lett.* **2004**, *6*, 2773.
- Roush, W. R.; Halvorsen, G. T. *Org. Lett.* **2007**, *9*, 2243.
- Maifeld, S. V.; Lee, D. *Org. Lett.* **2005**, *7*, 4995.
- Denmark, S. E.; Pan, W. T. *Org. Lett.* **2001**, *3*, 61.
- Hurley, A. L.; Welker, M. E.; Day, C. S. *J. Organomet. Chem.* **2000**, *598*, 150.
- Typical procedure for synthesis of siloxyenyne (**7e**). Alkenynol (**6e**)¹⁴ (1.236 g, 12.86 mmol), triethylamine (1.30 g, 12.85 mmol), dimethylaminopyridine (0.152 g, 1.267 mmol) and diisopropylchlorosilane (1.918 g, 12.73 mmol) were used to yield compound **7e** (1.680 g, 6.71 mmol, 53%) as a colorless, clear oily substance after purification by flash chromatography: *R_f* 0.85 (hexanes/Et₂O, 4:1); ¹H NMR (500 MHz, CDCl₃) δ 6.08 (h, *J* = 1.8 Hz, 1H, H-5), 4.48 (s, 2H, H-1), 4.20 (s, 2H, H-10), 1.98–2.17 (m, 4H, H6–9), 1.49–1.71 (m, 4H, H6–9), 0.96–1.12 (m, 14H, H-11, 12). ¹³C NMR (300 MHz, CDCl₃) δ 134.8 (C-5), 120.3 (C-4), 87.1 (C-2/3), 84.4 (C-2/3), 54.3 (C-1), 29.0 (C6–9), 25.6 (C6–9), 22.2 (C6–9), 21.5 (C6–9), 17.3 (C-12), 17.2 (C-12), 12.3 (C-11); HRMS calcd for C₁₅H₂₆NaOSi (M+Na)⁺ 273.1651, found 273.1639.
- General procedure for diisopropylsiloxacyclodiene synthesis (**8e**). (3-cyclohexenylprop-2-ynyl)diisopropylsilane (**7e**) (0.715 g, 2.854 mmol) was taken into a 50 mL, round bottom flask kept in a water bath at ambient temperature. The flask was purged with N₂ for 2 min, then 10 mol% KO^tBu (0.037 g, 0.330 mmol) was added in THF (3 × 5 mL) solution over a period of 10 min. After the addition, the water bath was removed and stirring continued for 1h at room temperature. The reaction mixture was diluted with Et₂O (20 mL), followed by quenching with satd. NH₄Cl (50 mL). The organic layer was separated and the aqueous layer was extracted with Et₂O (3 × 20 mL). The combined organics were washed with satd. NaCl solution (50 mL) and dried over MgSO₄. After removal of volatiles, the crude product was purified by flash chromatography yielding **8e** as a colorless, clear liquid; *R_f* 0.45 (hexanes/Et₂O, 15:2). The isolated compound was found to have an impurity of about ~20% with a similar *R_f* value, hence the compound was further purified by using a chromatotron (2.0 mm silica gel) (0.537 g, 2.144 mmol, 75%). ¹H NMR (300 MHz, CDCl₃) δ 6.56 (s, 1H, H-4), 5.57 (s, 1H, H-9), 4.61 (s, 1H, H-5), 2.17–2.26 (m, 2H, H-13), 2.03–2.16 (m, 2H, H-10), 1.64–1.75 (m, 2H, H-12), 1.52–1.64 (m, 2H, H-11), 1.06–1.20 (m, 2H, H-6), 1.04 (d, *J* = 6.6Hz, 6H, H-7), 0.98 (d, *J* = 6.8Hz, 6H, H-7); ¹³C NMR (300 MHz, CDCl₃) δ 140.3 (C-3), 138.0 (C-4), 135.7 (C-8), 128.9 (C-9), 77.8 (C-5), 26.2 (C-13), 26.0 (C-10), 22.8 (C-12), 22.4 (C-11), 17.4 (C-7), 17.1 (C-7), 13.4 (C-6); HRMS calcd for C₁₅H₂₇OSi (M+H)⁺ 251.1831, found 251.1828.
- Representative procedure for synthesis (tandem) of dimethylsiloxacyclodiene (**10a**): Alkenynol (**6a**) (1.792 g, 18.64 mmol) was taken into a 50 mL, round bottom flask kept in a water bath under N₂. After the slow addition of 1,1,3,3-tetramethyldisilazane (1.504 g, 11.28 mmol) over ~ 5 min using a syringe, the water bath (23 °C) was removed and stirring continued overnight at room temperature. Then volatiles were removed by rotovap (30 °C, 20mm) and the crude reaction mixture was dissolved in THF (10 mL), and the flask was cooled in a water-bath at ambient temperature. The flask was purged with N₂ for 2 min, then KO^tBu (0.217 g, 1.934 mmol) was added in THF (3 × 5 mL) solution over a period of 10 min. After the addition, the water bath was removed and stirring continued for 1h. at room temperature. The reaction mixture was diluted with Et₂O (20 mL), followed by quenching with satd. NH₄Cl (50 mL). The organic layer was separated and the aqueous layer was extracted with Et₂O (3 × 20 mL). The combined organics were washed with satd. NaCl solution (50 mL), dried over MgSO₄ and volatiles were removed by rotovap. A brown colored crude reaction mixture was produced which upon purification by column chromatography yielded **10a** as a clear colorless oil (2.014 g, 13.05 mmol, 70%); *R_f* 0.68 (pentane/Et₂O, 3:1); ¹H NMR (500 MHz, CDCl₃) δ 6.60 (as, 1H, H-4), 4.99 (s, 1H, H-8), 4.82 (s, 1H, H-8), 4.66 (s, 2H, H-5), 1.93 (s, 3H, H-9), 0.33 (as, 6H, H-6); ¹³C NMR (300 MHz, CDCl₃) δ 142.2 (C-3), 141.3 (C-7), 141.2 (C-4), 115.7 (C-8), 71.9 (C-5), 20.4 (C-9), 0.45 (C-6); HRMS calcd for C₈H₁₄OSi (M⁺) 154.0814, found 154.0813.